

OUTDOOR APPLICATION:
TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT): Use this product as a residual treatment to control Ticks, including the Deer Tick (*Ixodes dammini*), Lone Star Tick, (*Amblyomma americanum*), Brown Dog Tick (*Rhipicaphalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*). For each area listed below, apply until surface is wet (1/2 to 1 gallon for 750-1,000 sq. ft.)

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter. Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there. To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

CAMPING EQUIPMENT: MOSQUITO NETTING, NETS, TENTS, outsides of SLEEPING BAGS, BACKPACKS AND GROUND CLOTHS
For residual protection against ticks, chiggers and mosquitoes, treat camping equipment as follows:

SHAKE WELL BEFORE USING. Do not apply this product to clothing while it is being worn. Under NO circumstances should bare skin or clothing on the body be treated. Do not spray directly on or otherwise allow it to come in contact with skin, face or eyes. This product is to be used as a clothing and gear treatment only. Make all applications outdoors.

For six-week protection against ticks, chiggers and mosquitoes, select an outdoor area protected from the wind. Spray outer surfaces of clothing only (while not being worn) with a slow sweeping motion to lightly moisten the surface of the fabric, holding arm at a distance of 6 to 8 inches. (A complete outfit consists of shirt, trousers and socks.) Treat outside of each item of clothing for a minimum of 30 seconds on each side and allow 2 hours (4 hours under humid conditions) to dry prior to being worn. Pay particular attention to socks, trouser cuffs and shirt cuffs. Treatment should moisten the surface of the fabric enough to cause a slight color change. Use approximately 2 to 3 ounces of this container to treat the outsides of one complete set of clothing. Use remainder on gear. Follow normal laundering procedures weekly. Reapply after six weeks and sixth laundering.

In heavily infested areas, protection from ticks and chiggers will be improved by either wearing the cuffs inside footwear or taping pant cuffs around legs. Protection from mosquitoes can be further enhanced by applying an EPA approved repellent, as directed, to face, hands and other exposed areas of the body.

Do not treat inside of sleeping bag. Spray exterior surfaces of tent only. Select a well-ventilated outdoor area protected from wind and lay out gear to be treated. Hold container upright about 6 to 8 inches from surface of the article and spray with a slow sweeping motion to lightly moisten the surface of the fabric. Continue spraying over entire article until the outer surface of the article is moist enough to cause a light color change (darkening). Turn the article over and treat the other side. Hang treated gear and allow to dry for at least 2 hours (4 hours under humid conditions) before using. Do not retreat gear more than once every six weeks.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area out of reach of children and pets.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the container be used carefully following label directions and precautions.

CONTAINER HANDLING: IF EMPTY: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling, if available. IF PARTLY FILLED: Call your local waste agency or 1-800-253-2687 for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, absorb liquid with absorbent material such as soil or clay, etc. sweep up and dispose of material according to PRODUCT DISPOSAL above.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at **1-800-858-7378** or the poison control center at **1-800-222-1222**.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

PRECAUTIONARY STATEMENTS

Environmental Hazards
This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees and other pollinators exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees or other pollinators are foraging in the treatment areas.

Physical and Chemical Hazards
Do not use this product in or on electrical equipment due to the possibility of shock hazard. Do not allow product to freeze.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL BEFORE EACH USE.
FOR PET USE – USE ONLY ON DOGS**

USE RESTRICTIONS:

- Except when applying to fabric, dogs or horses, do not allow adults, children or pets to enter until vapors, mists and aerosols have dispersed and the treated area has been thoroughly ventilated.
- Except when applying to fabric, dogs or horses, do not allow adults, children or pets to enter the treated area until sprays have dried.
- Except when applying to fabric, dogs or horses, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not breathe vapors or spray mist.
- Do not apply when food is present.
- Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. Non-food/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, office, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).
- In the home, cover all food processing surfaces and utensils during treatment or thoroughly wash before use. Cover or remove exposed food.
- Do not contaminate food and feed stuffs. Remove pets, birds and cover fish aquaria before space spraying or surface applications.
- Not for broadcast use on indoor residential surfaces.
- Allow all treated articles to dry thoroughly before use.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawns, turf and other vegetation; 3) Applications to building foundations, up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.
- Do not water the treated area to the point of run-off.
- Do not make applications during rain.
- Application is prohibited directly into sewers or drains or to any area

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

SEE SIDE PANELS FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

**SHAKE WELL
BEFORE USE**



EPA REG. NO. 45385-99-56
EPA EST. NO. 35132-OH-1
NET CONTENTS: 1 Gallon / 3.785 Liters

ITEM #
209-W16



KILLS

BEDBUGS, TICKS & MOSQUITOES

REPELS MOSQUITOES THAT MAY CARRY AND TRANSMIT THE ZIKA VIRUS

PROTECTS TRAVEL LUGGAGE FROM "PICK-UP" OF BEDBUGS, AND OTHER LISTED INSECTS

FOR CONTROL OF TICKS WHICH MAY CARRY LYME DISEASE

ALSO FOR TREATMENT OF CLOTHING, TENTS AND CAMPING GEAR



Restrictions continued:
like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur. Do not allow product to enter any drain during or after application.

- Do not apply to classrooms while in use.
- Do not use in aircraft cabins.
- Not for use in outdoor residential misting systems.
- Not for use on meat or dairy animals, poultry or horses or foals intended for human consumption.
- NOT FOR USE ON HUMANS

SHAKE WELL BEFORE USING

Ready-to-Use and apply for most uses with the attached sprayer.

For travelers who want to protect their luggage from "picking up" of bed bugs, spiders, cockroaches and other listed insects, spray the outer and inner surfaces of luggage with a light spray mist of this product.

INDOOR APPLICATION

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING

TO KILL and CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, ANTS, SILVERFISH, SPIDERS AND CRICKETS: Apply at a rate of 1 gallon of product per 1000 sq. ft. (1 oz of product per 8 sq. ft.). Direct the spray into hiding places, cracks and crevices and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving.

TO CONTROL BED BUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floor. Repeat treatment as needed.

TO CONTROL FLEAS (ADULT AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE): Apply as a spot, crack & crevice treatment to infested areas of pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and floor and floor covering. Fresh bedding should be placed in animal quarters following treatment.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sun light, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying. Spray from a

distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry-cleaned.

SPACE SPRAYING

Close all doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths and Tobacco Moths: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1,000 cubic feet of space. Keep area closed for 2 hours.

INDOOR FOGGING: To kill exposed or accessible stages of the above named insects, use a rate of 1 quart to 1 gallon for 10,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours.

PEST CONTROL ON HORSES

To protect horses from: Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO KILL and CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: Apply at a rate of 2 ounces per adult horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the horse, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

FOR USE ON DOGS

To kill and control Fleas, Ticks and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat application every two weeks, if necessary. **DO NOT USE ON CATS OR KITTENS.**

Do not apply to dogs less than 12 weeks old. Consult a veterinarian before applying this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

EPA05292019/INV11202019 FNW209001JTE

Mfg. for J.T. EATON & CO., INC.,
1393 East Highland Road, Twinsburg, OH 44087 • U.S.A.
JT EATON™ is a trademark of J.T. EATON & CO., INC. ©2019
www.jteaton.com